

LAWNS

3 cups Top Choice Organic = 1 Pound

Established Lawns - Cool-Season (Northern) Grasses (Kentucky Bluegrass, Fescue, Perennial Ryegrass)

Application Date	Application Rate	Additional Information
Last week of May	2 lbs per 100 sq ft	Apply to dry grass only; water thoroughly after each application
First week of July	2 lbs per 100 sq ft	
First week of Sept	2 lbs per 100 sq ft	
Second week in Nov	2 lbs per 100 sq ft	Apply last fertilizer application mid November before first freeze. This is known as doormat feeding. This will provide an early spring green up.

Established Lawns - Warm-Season (Southern) Grasses (Bermuda, St. Augustine, Centipede grass, Bahia, & Zoysia)

Application Date	Application Rate	Additional Information
First week of April	2 lbs per 100 sq ft	Avoid nitrogen applications to these grasses less than 1 month than average first frost date in area. Bahia and Centipede grass require spring and summer applications. Avoid fertilization in late fall of these grasses to prevent winter kill
Last week of May	2 lbs per 100 sq ft	
First week of Sept	2 lbs per 100 sq ft	
Second week in Nov	2 lbs per 100 sq ft	

NEW Lawn

Application Date	Application Rate	Additional Information
Seed Bed	3 lbs per 100 sq ft	Rake lightly into soil, spread the new seed to recommended amounts, rake area lightly, roll and water the area thoroughly with a fine spray. Keep seed bed moist until seeds germinate.
Sodding prep	3 lbs per 100 sq ft	